

REC'D 25 JAN 2005

WIPO PCT


PA 1175924

THE UNITED STATES OF AMERICA**TO ALL TO WHOM THESE PRESENTS SHALL COME:****UNITED STATES DEPARTMENT OF COMMERCE****United States Patent and Trademark Office****May 26, 2004**

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A
FILING DATE UNDER 35 USC 111.**

APPLICATION NUMBER: 60/531,700**FILING DATE: December 22, 2003****PRIORITY
DOCUMENT****SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)**

**By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS**

**P. SWAIN****Certifying Officer**

PATENT APPLICATION SERIAL NO. _____

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICE
FEE RECORD SHEET

12/31/2003 EHAILE1 00000090 190134 60531700

01 FC:1005 160.00 DA

PTO-1556
(5/87)

*U.S. Government Printing Office: 2002 — 489-267/69033

17157 U.S. PTO
122203

Docket Number 4-33564P1

FILING BY "EXPRESS MAIL" UNDER 37 CFR 1.10

EV336719698US
Express Mail Label Number

December 22, 2003
Date of Deposit

Address to: **MS: Provisional Patent Application**
Commissioner for Patents
PO Box 1450
Alexandria, VA 22313-1450

19270 U.S. PTO
60/531700
122203

PATENT COVER SHEET FOR PROVISIONAL APPLICATION

Transmitted herewith for filing under 37 CFR §1.53(c) is the PROVISIONAL APPLICATION for patent of

INVENTOR(S)		
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
Anne Heidi Sauveur-Michel Terence	Boulay Lane Maira O'Reilly	Blotzheim, France Basel, Switzerland Habsheim, France Basel, Switzerland
TITLE OF THE INVENTION (280 characters max) ORGANIC COMPOUNDS		
CORRESPONDENCE ADDRESS Direct all correspondence to the address associated with Customer No. 001095, which is currently: Thomas Hoxie Novartis Corporate Intellectual Property One Health Plaza, Building 430 East Hanover, NJ 07936-1080		
ENCLOSED APPLICATION PARTS (check all that apply) <input checked="" type="checkbox"/> Specification (Including Any Claims and Abstract) - 10 pages <input type="checkbox"/> Drawings - sheets <input checked="" type="checkbox"/> Other (specify): Application Data Sheet		
METHOD OF PAYMENT The Commissioner is hereby authorized to charge filing fee and any additional fees, required to Deposit Account Number: 19-0134 in the name of Novartis.		
		PROVISIONAL FILING FEE AMOUNT: \$ 160

☐ U.S. Government agency and contract number: (if the invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.)

Respectfully submitted,

Thomas R. Savitsky

Thomas R. Savitsky
Attorney for Applicants
Reg. No. 31,661
Tel. No. (862) 778-7909

Date: December 22, 2003

INVENTOR INFORMATION

Inventor One Given Name:: Anne
Family Name:: Boulay
Postal Address Line One:: 9 rue des Pres
City:: Blotzheim
Country:: France
Postal or Zip Code:: F-68730
City of Residence:: Blotzheim
Country of Residence:: France
Citizenship Country:: France
Inventor Two Given Name:: Heidi
Family Name:: Lane
Postal Address Line One:: Lehenmattstrasse 189
City:: Basel
Country:: Switzerland
Postal or Zip Code:: 4052
City of Residence:: Basel
Country of Residence:: Switzerland
Citizenship Country:: Switzerland
Inventor Three Given Name:: Sauveur-Michel
Family Name:: Maira
Postal Address Line One:: 3c rue de Hombourg
City:: Habsheim
Country:: France
Postal or Zip Code:: F-68440 ,
City of Residence:: Habsheim
Country of Residence:: France
Citizenship Country:: France
Inventor Four Given Name:: Terence
Family Name:: O'Reilly
Postal Address Line One:: Drahtzugstrasse 51
City:: Basel
Country:: Switzerland
Postal or Zip Code:: 4057
City of Residence:: Basel
Country of Residence:: Switzerland
Citizenship Country:: Canada

CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 001095
Fax One:: 973-781-8064

APPLICATION INFORMATION

Title Line One:: ORGANIC COMPOUNDS
Total Drawing Sheets:: 0
Formal Drawings?: No

Application Type:: Provisional
Docket Number:: 4-33564P1
Secrecy Order in Parent Appl.?:: No

REPRESENTATIVE INFORMATION

Representative Customer Number:: 1095

Source:: PrintEFS Version 2.0

Organic Compounds

The present invention relates to biomarkers for determining the sensitivity of proliferative diseases such as cancer to therapeutic agents, in particular mTOR inhibitors.

A number of mTOR inhibitors have potent antiproliferative properties which make them useful for cancer chemotherapy, particularly of solid tumors, especially of advanced solid tumors. However there is still a need for more targeted use of mTOR inhibitors, which requires identification of patients which are likely to respond to treatment with such agents. Accordingly there is a need for biomarkers useful in e.g. clinical tests, which are capable of predicting responsiveness of a proliferative disease, e.g. a tumor in a patient to treatment with an mTOR inhibitor.

The present invention provides in one aspect use of S6 40S ribosomal protein (S6), in particular phosphorylated S6, as a biomarker for determining the sensitivity of a proliferative disease to treatment with an mTOR inhibitor.


In a further aspect the invention provides a method for determining the sensitivity of a proliferative disease in a subject to treatment with an mTOR inhibitor, comprising determining the level of expression and/or phosphorylation state of S6 in a sample derived from the subject.

In another aspect the invention provides a method of selecting subjects suffering from a proliferative disease for treatment with an mTOR inhibitor, comprising determining the sensitivity of the proliferative disease to treatment with an mTOR inhibitor in each subject by a method as described above, and selecting those subjects showing increased expression of phosphorylated S6 for treatment with an mTOR inhibitor.

It has surprisingly been found that S6 40S ribosomal protein (otherwise known as S6) is a useful biomarker which is predictive of sensitivity of proliferative diseases to treatment with an mTOR inhibitor. In particular, it has been found that the phosphorylation state of S6 correlates well with sensitivity to mTOR inhibitors. mTOR inhibitors are more likely to show a significant antiproliferative effect when used to treat cancer cell lines showing higher levels of expression of phosphorylated S6. S6 is a component of the 40S ribosomal subunit which

is a substrate for the p70 S6 kinase, a downstream effector of the mTOR protein kinase. Multiple phosphorylation of S6 has been implicated in the translational up-regulation of mRNAs encoding components of the protein synthetic apparatus, and as such is thought to play a major role in the growth of mammalian cells (Volarevic and Thomas, Prog. Nucleic Acid Res. Mol. Biol. 2001, 65:101-27). The sequence of human S6 is available under Genbank accession number M20020.

The term "mTOR inhibitor" as used herein includes, but is not limited to rapamycin (sirolimus) or a derivative thereof. Rapamycin is a known macrolide antibiotic produced by *Streptomyces hygroscopicus*. Suitable derivatives of rapamycin include e.g. compounds of formula A



A

wherein

R1aa is CH₃ or C₃₋₆alkynyl,

R2aa is H or -CH₂-CH₂-OH, 3-hydroxy-2-(hydroxymethyl)-2-methyl-propanoyl or tetrazolyl, and

Xaa is =O, (H,H) or (H,OH)

provided that R2aa is other than H when Xaa is =O and R1aa is CH₃.

or a prodrug thereof when R2aa is -CH₂-CH₂-OH, e.g. a physiologically hydrolysable ether thereof.

Compounds of formula A are disclosed e.g. in WO 94/09010, WO 95/16691, WO 96/41807, USP 5,362,718 or WO 99/15530 which are incorporated herein by reference. They may be prepared as disclosed or by analogy to the procedures described in these references.

Preferred rapamycin derivatives are 32-deoxorapamycin, 16-pent-2-ynyloxy-32-deoxorapamycin, 16-pent-2-ynyloxy-32(S)-dihydro-rapamycin, 16-pent-2-ynyloxy-32(S)-dihydro-40-O-(2-hydroxyethyl)-rapamycin and, more preferably, 40-O-(2-hydroxyethyl) rapamycin. Further examples of rapamycin derivatives include e.g. CCI779 or 40- [3-hydroxy-2-(hydroxymethyl)-2-methylpropanoate]-rapamycin or a pharmaceutically acceptable salt thereof, as disclosed in USP 5,362,718, ABT578 or 40-(tetrazolyl)-rapamycin, particularly 40-epi-(tetrazolyl)-rapamycin, e.g. as disclosed in WO 99/15530, or rapalogs as disclosed e.g. in WO 98/02441 and WO01/14387, e.g. AP23573 or TAFA-93.

In each case where citations of patent applications or scientific publications are given, the subject-matter relating to the compounds is hereby incorporated into the present application by reference. Comprised are likewise the pharmaceutically acceptable salts thereof, the corresponding racemates, diastereoisomers, enantiomers, tautomers as well as the corresponding crystal modifications of above disclosed compounds where present, e.g. solvates, hydrates and polymorphs, which are disclosed therein. The compounds used as active ingredients in the combinations of the invention can be prepared and administered as described in the cited documents, respectively.

The proliferative disease may be a neoplastic disease, preferably a malignant proliferative disease, e.g. a cancer, e.g. a solid tumor, particularly an advanced solid tumor as disclosed in WO 02/66019. By "solid tumors" are meant tumors and/or metastasis (wherever located) other than lymphatic cancer, e.g. brain and other central-nervous system tumors (eg. tumors of the meninges, brain, spinal cord, cranial nerves and other parts of central nervous system, e.g. glioblastomas or medulla blastomas); head and/or neck cancer; breast tumors; circulatory system tumors (e.g. heart, mediastinum and pleura, and other intrathoracic organs, vascular tumors and tumor-associated vascular tissue); excretory system tumors (e.g. kidney; renal pelvis, ureter, bladder, other and unspecified urinary organs); gastrointestinal tract tumors (e.g. oesophagus, stomach, small intestine, colon, colorectal, rectosigmoid junction, rectum, anus and anal canal), tumors involving the liver and intrahepatic bile ducts, gall bladder, other and unspecified parts of biliary tract, pancreas, other and digestive organs); head and neck; oral cavity (lip, tongue, gum, floor of mouth, palate, and other parts of mouth, parotid gland, and other parts of the salivary glands, tonsil,

oropharynx, nasopharynx, pyriform sinus, hypopharynx, and other sites in the lip, oral cavity and pharynx); reproductive system tumors (e.g. vulva, vagina, Cervix uteri, Corpus uteri, uterus, ovary, and other sites associated with female genital organs, placenta, penis, prostate, testis, and other sites associated with male genital organs); respiratory tract tumors (e.g. nasal cavity and middle ear, accessory sinuses, larynx, trachea, bronchus and lung, e.g. small cell lung cancer or non-small cell lung cancer); skeletal system tumors (e.g. bone and articular cartilage of limbs, bone articular cartilage and other sites); skin tumors (e.g. malignant melanoma of the skin, non-melanoma skin cancer, basal cell carcinoma of skin, squamous cell carcinoma of skin, mesothelioma, Kaposi's sarcoma); and tumors involving other tissues including peripheral nerves and autonomic nervous system, connective and soft tissue, retroperitoneum and peritoneum, eye and adnexa, thyroid, adrenal gland and other endocrine glands and related structures, secondary and unspecified malignant neoplasm of lymph nodes, secondary malignant neoplasm of respiratory and digestive systems and secondary malignant neoplasm of other sites. Where hereinbefore and subsequently a tumor, a tumor disease, a carcinoma or a cancer is mentioned, also metastasis in the original organ or tissue and/or in any other location are implied alternatively or in addition, whatever the location of the tumor and/or metastasis is.

According to the method of the present invention, subjects suffering from such a proliferative disease can be screened in order to predict their sensitivity to mTOR inhibitors. The method may be performed in vitro, e.g. on a sample of biological tissue derived from the subject. The sample may be any biological material separated from the mammalian body such as e.g. tissue, cell lines, plasma or serum, cell or tissue lysate, preferably tumor tissue. The subject is preferably a human subject.

Levels of expression and/or phosphorylation state of S6 are assayed in the biological sample by any technical means on the basis of e.g. RNA expression using for example the technique of RT-PCR or on the basis of e.g. protein expression using for example the technique of Western blotting, immunohistochemistry or ELISA, including immunoassays, immunoprecipitation and electrophoresis assays. Preferably the method comprises determining the level of expression of (e.g. human) S6 protein, and in particular phosphorylated S6 in the sample. The method may involve detection of phosphorylation at any phosphorylation site on S6. For example, phosphorylation of (e.g. human) S6 on serine

235/236 may be determined, more preferably phosphorylation of S6 on serines 240/244 is determined.

For example, antibodies specific for (e.g. phosphorylated) S6 are used in a standard immunoassay format to measure (e.g. phosphorylated) S6 levels. ELISA (enzyme linked immunosorbent assay) type assays, immunoprecipitation type assays, conventional Western blotting assays and immunohistochemistry assays using e.g. monoclonal or polyclonal antibodies are also utilized to determine levels of the phosphorylated S6 as a biomarker protein.

Polyclonal and monoclonal antibodies specific to S6, e.g. to S6 protein or to phosphorylated S6 are produced in accordance with known immunization methods.

The phosphorylated S6 level may also be measured by two-dimensional (2-D) gel electrophoresis. 2-D gel electrophoresis is known in the art and typically involves isoelectric focusing (IEF) along a first dimension followed by SDS-PAGE (sodium dodecyl sulphate-polyacrylamide gel electrophoresis) along a second dimension. The resulting electropherograms are analyzed, for example, by immunoblot analysis using antibodies. Suitable antibodies directed against S6 protein or phosphorylated S6 can be produced as discussed above or obtained from a commercial source (e.g. Cell Signaling Technology® catalogue # 2212; #2215; #2211).

The present invention thus provides a method of screening subjects suffering from a proliferative disease in order to predict their responsiveness to treatment with an mTOR inhibitor, comprising determining the level of expression and/or phosphorylation state of S6 by a method as defined above.

In a further aspect, the present invention provides a method of treating a proliferative disease in a subject in need thereof, comprising determining the level of expression and/or phosphorylation state of S6 in a sample derived from the subject, by a method as described above, and treating the subject with an mTOR inhibitor if the level of expression of (e.g. phosphorylated) S6 is elevated.

The level found in a particular tissue from a subject, e.g. a sample of tumor tissue, may be compared with a control sample, e.g. a sample of normal tissue from a subject not suffering from the disease, or a sample of normal (i.e. non-tumor) tissue from the same subject. An elevated level of phosphorylated S6, e.g. above control levels, is predictive of a beneficial therapeutic effect (i.e. an antiproliferative effect) of an mTOR inhibitor. The elevated level at which use of an mTOR inhibitor is indicated may be determined by a skilled person, e.g. in certain embodiments treatment with an mTOR inhibitor may be indicated where the level of phosphorylated S6 in the sample is detectably above the control level, or where the level is at least 50%, 100%, 500% or 1000% higher than control.

Moreover, the method may be used to select an appropriate dose of an mTOR inhibitor in order to individually optimise therapy for each patient. For instance a lower dose of an mTOR inhibitor may be selected where a sample from the subject shows higher phosphor-S6 levels, and vice versa. Factors for consideration in this context include the particular condition being treated, the particular mammal being treated, the clinical condition of the individual patient, the site of delivery of the active compound, the particular type of the active compound, the method of administration, the scheduling of administration, the severity of the condition and other factors known to medical practitioners. The therapeutically effective amount of an active compound to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat the disease. Such amount is preferably below the amount that is toxic to the host or which renders the host significantly more susceptible to infections. Appropriate doses of an mTOR inhibitor are e.g. as disclosed in WO 02/66019, e.g. daily dosage rates of the order of ca. 0.1 to 25 mg, e.g. from ca. 0.05 to 10 mg active ingredient p.o., as a single dose or in divided doses or intermittent, e.g. once a week. Rapamycin or a derivative thereof, e.g. a compound of formula A, may be administered by any conventional route, in particular enterally, e.g. orally, e.g. in the form of tablets, capsules, drink solutions or parenterally, e.g. in the form of injectable solutions or suspensions, containing, for example, from about 0.1 % to about 99.9%, preferably from about 1 % to about 60 %, of the active ingredient(s).

Example 1

Human tumor cell lines, e.g. the 40-O-(2-hydroxyethyl) rapamycin-sensitive MCF7, BT549 or LNCap lines (IC50 in sub nM range) versus the comparative 40-O-(2-hydroxyethyl)

rapamycin-resistant PC3M line (IC_{50} in the > 100 nM range), as well as cell lines with moderate rapamycin-sensitivity ($1 \text{ nM} < IC_{50} < 100 \text{ nM}$) such as DU145, HCC1937 and MDA-MB231, are added to 96-well plates (1,000 to 5000 cells/well in 100 μ l medium) and incubated for 24 hr. Subsequently, a dilution series of an mTOR inhibitor, e.g. a compound of formula A, e.g. 40-O-(2-hydroxyethyl) rapamycin is made in separate wells and the dilutions are added to the wells. The cells are then re-incubated for 4 days. Methylene blue staining is performed on day 5 and the amount of bound dye (proportional to the number of surviving cells that bind the dye) determined. IC_{50} s are subsequently determined using Softmax 1.2.0 software.

The same tumor cell lines as above, cultured to 50 – 70 % confluency, are refed with normal culture medium (10 % v/v FCS). Protein lysates are prepared and 20 μ g electrophoretically resolved and transferred to polyvinylidene difluoride (PVDF) by semi-dry electroblotting. Blots are probed with anti-phospho-S6 or anti-S6 protein antibody and decorated proteins are revealed using enhanced chemiluminescence. The relative intensities of S6 phosphorylation in each cell line are revealed and numerated as: 0 (no phosphorylation observed), 0.5, 1, 2, 3 or 4 (Maximal phosphorylation observed).

Comparison of phosphorylated S6 levels with IC_{50} measurements for the mTOR inhibitor in the same cell lines indicate a significant correlation between increased antiproliferative activity of the mTOR inhibitor and increased levels of phosphorylated S6 ($n = 7$, $R = -0.746$, $p = 0.00384$ by Spearman Rank correlation analysis). A similar correlation was not observed when performing the same analysis with phosphorylated MAPK (ERK1/2; $n = 8$, $R = -0.0755$, $p = 0.839$).

In order to predict sensitivity of e.g. a tumor in a subject to mTOR inhibitors, a similar analysis to that described above is performed using a sample containing tumor tissue from the subject in place of the human tumor cell lines. Phosphorylated S6 levels obtained from the tumor tissue sample may be compared with that obtained from control tissue, or with data obtained from the human tumor cell lines, in order to predict likely responsiveness to an mTOR inhibitor.

CLAIMS

1. Use of S6 as a biomarker for determining the sensitivity of a proliferative disease in a subject to treatment with an mTOR inhibitor.
2. Use according to claim 1, comprising use of the level of expression and/or phosphorylation state of S6.
3. Use according to any preceding claim, comprising use of the level of expression of phosphorylated S6 protein.
4. A method for determining the sensitivity of a proliferative disease in a subject to treatment with an mTOR inhibitor, comprising determining the level of expression and/or phosphorylation state of S6 in a sample derived from the subject.
5. A method or use according to any preceding claim, wherein the proliferative disease comprises a cancer.
6. A method or use according to any preceding claim, wherein the mTOR inhibitor comprises rapamycin or a rapamycin derivative.
7. A method or use according to claim 6, wherein the rapamycin derivative comprises 40-O-(2-hydroxyethyl) rapamycin.
8. A method according to any of claims 3 to 7, comprising determining the level of expression of phosphorylated S6 protein.
9. A method according to any of claims 3 to 8, wherein the sample is derived from a tumor in the subject.
10. A method according to any of claims 3 to 9, wherein increased expression of phosphorylated S6 relative to control is predictive of sensitivity of the proliferative disease to treatment with the mTOR inhibitor.

11. A method of selecting subjects suffering from a proliferative disease for treatment with an mTOR inhibitor, comprising determining the sensitivity of the proliferative disease to treatment with an mTOR inhibitor in each subject by a method as described in any of claims 3 to 10, and selecting those subjects showing increased expression of phosphorylated S6 for treatment with an mTOR inhibitor.

12. A method of treating a proliferative disease in a subject in need thereof, comprising determining the level of expression of phosphorylated S6 in a sample derived from the subject, by a method as described in any of claims 3 to 10, and treating the subject with an mTOR inhibitor if the level of expression of phosphorylated S6 is elevated.

ABSTRACT

Disclosed is a method for determining the sensitivity of a proliferative disease in a subject to treatment with an mTOR inhibitor, comprising determining the level of expression and/or phosphorylation state of S6 in a sample derived from the subject, as well as related methods of treatment and uses.